

Dynamical Magnetic Susceptibility for the t - J Model

Th. Pruschke, Th. Obermeier and J. Keller

Institut für Festkörperphysik, Universität Regensburg, 93040 Regensburg, Germany

We present results for the *dynamical* magnetic susceptibility of the t - J model, calculated with the dynamical mean field theory. For $J = 0$ we find enhanced ferromagnetic correlations but an otherwise relatively $\vec{q}$ -independent dynamical magnetic susceptibility. For $J > 0$ the explicit antiferromagnetic exchange leads to a dynamic spin structure factor with the expected peak at the antiferromagnetic Bragg point.

The observation that the local electron-electron interaction in models of strongly correlated electrons leads in the limit spatial dimensions $d \rightarrow \infty$ to a local one-particle self energy [1], founded the so-called *dynamical mean field theory* (DMFT): The lattice problem maps onto an impurity system coupled to an effective, self-consistently determined bath [2], the lattice structure enters only via the free density of states (DOS) $\rho_0(\epsilon) = \sum_{\vec{k}} \delta(\epsilon - \epsilon_{\vec{k}})$.

In addition to the calculation of one-particle properties the DMFT also allows to obtain the two-particle quantities [3], especially the *dynamical* magnetic susceptibility. So far, however, only static susceptibilities have been studied [3, 4, 5]. In this contribution we want to present first results for the dynamical magnetic susceptibility of a strongly correlated electron system calculated within the DMFT.

The model we want to discuss is the well-known t - J model

$$H_{tJ} = -t \sum_{\langle i,j \rangle \sigma} \hat{c}_{j\sigma}^\dagger \hat{c}_{i\sigma} + J \sum_{\langle i,j \rangle} \vec{S}_i \cdot \vec{S}_j .$$

The operator $\hat{c}_{i\sigma}^\dagger$ creates an electron with spin σ on site i if and only if no other particle is present, the $\vec{S}_i$ are the conventional spin operators and $\langle \cdot, \cdot \rangle$ denotes sums on nearest neighbours only. In a previous publication [5], we have shown that the magnetic susceptibility for this model acquires a particularly simple form, namely $\chi_{\vec{q}}^{tJ}(z) = \chi_{\vec{q}}^\infty(z) \left[1 - J_{\vec{q}} \chi_{\vec{q}}^\infty(z) \right]^{-1}$. Here, $\chi_{\vec{q}}^\infty(z)$ denotes the magnetic susceptibility for $J = 0$, i.e. the $U = \infty$ Hubbard model, and $J_{\vec{q}}$ is given by

$$J_{\vec{q}} = -\frac{J}{d} \sum_{l=1}^d \cos(q_l \cdot a) .$$

The major problem is the calculation of $\chi_{\vec{q}}^\infty(z)$. This task can be performed most conveniently in Matsubara space, since there the calculation reduces to the inversion of matrices. The final step, the analytic continuation $\chi_{\vec{q}}^\infty(i\nu_n) \rightarrow \chi_{\vec{q}}^\infty(\omega + i\delta)$ is then performed with Padé approximation. To calculate $\chi_{\vec{q}}^\infty(z)$ we use standard resolvent techniques in connection with the NCA [6] as explained in reference [5]. Note that since we solve




Figure 1: Local dynamical magnetic susceptibility for $J = 0$ obtained directly from the effective local model (lines) and from the Matsubara data via numerical analytic continuation with Padé approximants (symbols). The inset shows the underlying data in Matsubara space. The parameters in the calculation are $\delta = 5\%$ and $T = t/30$.

the DMFT by perturbative techniques the results for $\chi_{\vec{q}}^\infty(i\nu_n)$ do not contain statistical errors like

in QMC and thus the analytic continuation via Padé approximants is expected to work properly. Let us nevertheless start with some “confidence tests” for the latter procedure. This can most conveniently be done with the help of the local susceptibility, where one can obtain both, $\chi_{loc}(\omega + i\delta)$ and $\chi_{loc}(i\nu_l)$ *independently*. In Fig. 1 we compare directly calculated results for the local dynamic susceptibility (full lines) to those obtained from a Padé approximation (symbols). The typically rather structureless data on the imaginary axis are displayed in the inset. Note that we used merely $n = 64$ Matsubara frequencies in the Padé approximation. This example shows that at least for mildly varying functions like susceptibilities the Padé approximation is a rather accurate tool for numerical analytic continuation.




Figure 2: Spin structure factor at $J = 0$ for $\delta = 5\%$ and $T = t/30$ as function of $\vec{q}$ and ω . Most prominent is a slight enhancement of *ferromagnetic* fluctuations for $\omega \rightarrow 0$ but an otherwise rather weak variation with $\vec{q}$.

The complete dynamical spin-structure factor ($g(\omega)$ denotes the Bose function) $S(\vec{q}, \omega) = \Im m \left\{ \chi_{\vec{q}}^{\infty}(\omega + i0^+) \right\} \cdot \{1 + g(\omega)\}$ for $J = 0$, i.e. the $U = \infty$ Hubbard model, is shown in Fig. 2. The doping is $\delta = 5\%$ and the temperature $T = t/30$.

The result is quite characteristic for the spin correlations in this model: As in the static case [5], we find a rather weak $\vec{q}$ -dependence of $S(\vec{q}, \omega)$. An interesting aspect is that for $\omega \rightarrow 0$ the ferromagnetic correlations at $q = 0$ are enhanced, while for increasing energy $\vec{q} = (\pi, \pi, \dots)$ slightly dominates. This result can be understood by the fact that, since there is no net magnetic exchange present in this model, no special configuration is preferred by the system, except for $\omega \rightarrow 0$ where a small gain in kinetic energy by a partially ferromagnetic polarization [7] is responsible for enhanced ferromagnetic correlations.




Figure 3: Spin structure factor for the t - J model with $J = T$, other parameters as in Fig. 2. The fluctuations are now strongest for $\omega \rightarrow 0$ and $\vec{q}$ at the antiferromagnetic Bragg point, as expected.

The situation of course changes when we include a finite exchange $J = T$: As is clear from Fig. 3, the antiferromagnetic correlations are strongly enhanced, as expected. It is noteworthy that the weak $\vec{q}$ -dependence in $\chi_{\vec{q}}^{\infty}(\omega + i0^+)$ has no significant influence for larger J . Note that the value of J chosen here is still only 50% of the critical value where a transition into an ordered state occurs for the current parameter values. This supports our earlier suggestion that the prominent effects of the spin dynamics for the t - J model in the paramagnetic phase can already be addressed

by replacing $\chi_{\vec{q}}^{\infty}(\omega + i0^+) \rightarrow \chi_{loc}(\omega + i0^+)$ [5]. To conclude we presented DMFT-results for the dynamical susceptibility for the t - J model. We showed that for the present problem the numerical analytic continuation by means of Padé approximation is a reliable tool. The results for the dynamical susceptibility of the $U = \infty$ Hubbard model show a behaviour as already conjectured from the statics [5], namely a rather weak variation with $\vec{q}$ and a slight enhancement of the ferromagnetic correlations as $\omega \rightarrow 0$. A finite exchange interaction J again strongly enhances the antiferromagnetic fluctuations. In this case the detailed $\vec{q}$ structure of the underlying susceptibility of the Hubbard model is completely washed out. This observation gives to some extent a microscopic justification of recent suggestions for approximate forms of the dynamical magnetic susceptibility for strongly correlated models [8].

The authors would like to thank Dr. A. Liechtenstein for providing the code for the Padé approximations. This work was supported by the Deutsche Forschungsgemeinschaft grant number Pr 298/3-1.

References

- [1] W. Metzner and D. Vollhardt, Phys. Rev. Lett. **62**, 324 (1989); E. Müller-Hartmann, Z. Phys. B **74**, 507 (1989).
- [2] V. Janiš and D. Vollhardt, Int. J. Mod. Phys. B **6**, 731(1992); M. Jarrell, Phys. Rev. Lett. **69**, 168(1992); A. Georges and G. Kotliar, Phys. Rev. **B45**, 6479(1992).
- [3] M. Jarrell and Th. Pruschke, Z. Phys. B. **90**, 187(1993).
- [4] J.K. Freericks and M. Jarrell, Phys. REv. Lett. **74**, 186(1995).
- [5] Th. Pruschke, Q. Qin, Th. Obermeier and J. Keller, J. Phys.: Condens. Matter **8**, 3161(1996).
- [6] N. E. Bickers, D. L. Cox, J. W. Wilkins, Phys. Rev. **B36**, 2036(1987).
- [7] Y. Nagaoka, Phys. Rev. **147**, 392(1966).
- [8] P. Monthoux und D. Pines, Phys. Rev. **B49**, 4261(1994); P. Monthoux und D.J. Scalapino, Phys. Rev. Lett. **72**, 1874(1994).